



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

NUMBER THEORY.

When this issue was made up no solutions had been received for numbers 189, 191, 192, 196, 201, and 202. Please give attention to these.

208. Proposed by E. T. BELL, Seattle, Washington.

If an odd number is perfect, it cannot be the sum of two squares.

209. Proposed by R. D. CARMICHAEL, Indiana University.

Prove that the difference of the sixth powers of two integers cannot be the square of an integer

210. Proposed by ELMER SCHUYLER, Brooklyn, N. Y.

If a and b are relatively prime and $(a + b)$ is even, then $(a - b)ab(a + b) \equiv 0 \pmod{24}$.

SOLUTIONS OF PROBLEMS.

ALGEBRA.

394. Proposed by E. B. ESCOTT, Ann Arbor, Michigan.

Solve the equation $\sin^2 x \sin^2 2x = 5/16$.

SOLUTION BY R. M. MATHEWS, Riverside, California.

Taking the square root of each side of the given equation we obtain

$$(1) \sin x \sin 2x = \frac{\sqrt{5}}{4} \quad \text{and} \quad (2) \sin x \sin 2x = -\frac{\sqrt{5}}{4}.$$

From (1),

$$2 \sin^2 x \cos x = \frac{\sqrt{5}}{4}, \quad \text{or} \quad (3) \cos^3 x - \cos x + \frac{\sqrt{5}}{8} = 0.$$

Inspection shows that $-\frac{\sqrt{5}}{2}$ is a value of $\cos x$ in (3). Eliminating this impossible solution, we have the quadratic

$$(4) \cos^2 x - \frac{\sqrt{5}}{2} \cos x + \frac{1}{4} = 0,$$

the roots of which are $\cos x = \frac{\sqrt{5} + 1}{4}$ and $\cos x = \frac{\sqrt{5} - 1}{4}$.

The solutions of (2) are the negatives of those of (1) just found.

The side of a regular inscribed decagon is $(\sqrt{5} - 1)/2$, whence $(\sqrt{5} - 1)/4 = \cos 72^\circ$. From this, it is easily shown that $(\sqrt{5} + 1)/4 = \cos 36^\circ$.

Since in the original equations the functions are all squared, the signs which they may have in particular quadrants are immaterial. Hence

$$x = n\pi \pm \frac{1}{5}\pi, \quad x = n\pi \pm \frac{2}{5}\pi,$$

where n is any positive or negative integer, are the real values satisfying the equation.

Also solved by C. E. GITHENS, HORACE OLSON, ALBERT R. NAUER, RICHARD MORRIS, A. M. HARDING, ELMER SCHUYLER, S. W. REAVES, W. C. EELLS, LEROY M. COFFIN, A. H. HOLMES, F. M. MORGAN, H. C. FEEMSTER, J. L. RILEY, H. E. TREFETHEN, and the PROPOSER.

395. Proposed by V. M. SPUNAR, Chicago, Illinois.

Solve the system of equations

$$x_1^2 x_2 = a_1, \quad x_2^2 x_3 = a_2, \quad x_3^2 x_4 = a_3, \quad \dots, \quad x_n^2 x_1 = a_n.$$